



In the claims:

Claim 1 (currently amended) A Process process for the synthesis of a ceramide-type compounds, characterized in that it includes at least an amide formation step, performed by means of the lipase B-type enzyme of *Candida antartica*, and an esterification step, also performed by means of a lipase-type enzyme, and in that the ceramide-type compounds correspond to the general formula (1):



in which the group Am-Alc figures a C2 to C6 carbon chain, preferably saturated, linear or optionally branched, obtained from an amino alcohol; X figures a hydrogen atom or a C1 to C4 carbon chain, optionally hydroxylated on the 2' and/or following positions of the amino group; and in which each of the groups AG and AG' figures a C4 to C30 carbon chain, saturated or unsaturated, obtained from a fatty acid or a fatty acid ester; the two groups AG and AG' may be identical or different of the formula



wherein Am-Alc is alkyl of 2 to 6 carbon atoms derived from an amino alcohol, X is hydrogen or alkyl of 1 to 4 carbon atoms optionally hydroxylated in the 2' position

and AG and AG' are individually unsaturated or saturated hydrocarbon of 4 to 30 carbon atoms derived from a fatty acid or fatty acid amide comprising reacting on amino alcohol of the formula



with an acid of the formula AG-COOH wherein AG is defined as above in the presence of a lipase B-type enzyme of *Candida antartica* to introduce the

AG-C-N- group and with an AG'-COOH wherein AG' is defined as above in the presence of *Rhizomucor miehei* lipase to introduce the AG'-C-O group.

Claim 2 (currently amended) ~~Process according to~~ The process of
 claim 1, ~~characterized in that~~ wherein the amide formation step is carried out under stoichiometric conditions between a the fatty acid ~~and/or its ester and an~~ or the amino-alcohol at a temperature comprised ~~between~~ of 40 and to 100°C.

Claim 3 (currently amended) ~~Process according to~~ The process of
 claim 1, ~~characterized in that~~ wherein the amide formation is carried out without solvent, at a ~~minimal~~ temperature of about 65°C.

Claim 4 (currently amended) ~~Process according to~~ The process of
 claim 1, ~~characterized in that~~ wherein the amide formation is carried out under a

reduced pressure ~~comprised between~~ of 1 and to 500 mbars ~~and~~ during at least 16 hours.

Cancel Claim 5.

Claim 6 (currently amended) ~~Process according to~~ The process of
claim 5 1, ~~characterized in that~~ wherein the esterification reaction is carried out with
a ratio fatty acid ester/ to amino-alcohol ~~comprised between~~ of 1 and to 2.

Claim 7 (currently amended) ~~Process according to~~ The process of
claim 5 1, ~~characterized in that~~ wherein the esterification reaction is carried out at a
temperature ~~comprised between~~ of 40 and to 90°C.

Claim 8 (currently amended) ~~Process according to~~ The process of
claim 5 1, ~~characterized in that~~ wherein the esterification reaction is carried out
without solvent, at a ~~minimal~~ temperature of about 65°C.


Claim 9 (currently amended) ~~Process according to~~ The process of
claim 5 1, ~~characterized in that~~ wherein the esterification reaction is carried out
under a reduced pressure ~~comprised between~~ of 1 and to 500 mbars ~~and~~ during at
least 18 hours.

Claim 10 (currently amended) ~~Process according to~~ **The process of**

Claim 11 (currently amended) ~~Process according to~~ The process of

Cancel Claim 12.

Claim 13 (currently amended) ~~Process according to~~ The process of



in which:

~~-n is an integer selected from the numbers 1, 2, or 3 and m is an integer selected from the numbers 1, 2, or 3,~~

~~-X is selected from the group composed of hydrogen and a C1 to C4 carbon chain, optionally hydroxylated on the positions 2' and/or followings of the amino group;~~

~~-R¹ is selected from the group composed of hydrogen and a C1 to C4 carbon chain, preferably saturated, linear, optionally branched and/or hydroxylated or alkyl of 1 to 4 carbon atoms optionally hydroxylated,~~

~~-R² is selected from the group composed consisting of hydrogen, -OH, -NH₂ and a C1 to C4 carbon chain alkyl of 1 to 4 carbon atoms, preferably saturated, linear, optionally branched and/or hydroxylated,~~

~~-R³ is selected from the group composed consisting of hydrogen, -OH, -CH₂OH, and in which at least one of the groups R¹, R² and R³ includes a -OH group.~~

Claim 14 (currently amended) ~~Process according to~~ The process of
claim 1 ~~characterized in that~~ wherein the amide formation step is performed before
the esterification step.